











Marine Carotenoids

Takashi Maoka, Ph.D.

Research Institute for Production Development, Sakyo-ku, Kyoto, Japan 606-0805
(maoka@mbox.kyoto-inet.or.jp)

New carotenoids from Oyster *Crassostrea gigas*




Yasuhiro Fujiwara, Takashi Maoka, Masahiro Ookubo and Takao Matsuno, Crassostreaxanthin A and B, Novel marine carotenoids from the oyster *Crassostrea gigas*, *Tetrahedron Lett*, **33**, 4941-4944 (1992).

Takashi Maoka, Keiji Hashimoto, Naoshige Akimoto and Yasuhiro Fujiwara, Structures of Five New Carotenoids from the Oyster *Crassostrea gigas*, *J. Nat. Prod.*, **64**, 578-581 (2001).


Takashi Maoka, Yasuhiro Fujiwara, Keiji Hashimoto, and Naoshige Akimoto, Structures of New Carotenoids with a 3,4-Dihydroxy- β -End Group from the Oyster *Crassostrea gigas*, *Chem. Pharm. Bull.*, **53**, 1207-1209 (2005).

Chinese Surf Clam, *Macra chinensis*



Takashi Maoka, Yasuhiro Fujiwara, Keiji Hashimoto, and Naoshige Akimoto, Characterization of Fucoxanthin and Fucoxanthinol Esters in the Chinese Surf Clam, *Macra chinensis*, *J. Agric. Food Chem.*, **55**, 1563 – 1567 (2007).


Corbicula clam *Corbicula japonica*



Takashi Maoka, Yasuhiro Fujiwara, Keiji Hashimoto, and Naoshige Akimoto, Structure of New Carotenoids from the Corbicula clam *Corbicula japonica*, *J. Nat. Prod.*, **68**, 1341-1344 (2005).


Takashi Maoka, Yasuhiro Fujiwara, Keiji Hashimoto, and Naoshige Akimoto, Carotenoids in Three Species of Corbicula Clams, *Corbicula japonica*, *Corbicula sandai*, and *Corbicula* sp. (Chinese Freshwater Corbicula Clam), *J. Agric. Food Chem.*, **53**, 8357-8364 (2005).

***Paphia amabilis* (Satuma Akagai in Japanese)**



Takashi Maoka, Naoshige Akimoto, Mi-Jin Yim, Masashi Hosokawa, Kazuo Miyashita, A New C₃₇-Skeletal Carotenoid from the Clam, *Paphia amabilis*. *J. Agric Food Chem.*, **56**, 12069-12072 (2008).


Spindle shell *Fushinus perplexus*



Takao Matsuno, Kiyoshi Katagiri, Takashi Maoka and Tadaaki Komori, Novel reductive metabolic pathways of 4-oxo- β -end group in carotenoids of the spindle shell *Fushinus perplexus*, *Comp. Biochem. Physiol.*, **81B**, 905-908 (1985).

Miyuki Tsushima, Takashi Maoka and Takao Matsuno, Structure of Carotenoids with 5,6-Dihydro- β -End Groups from the Spindle Shell *Fushinus perplexus*, *J. Nat. Prod.*, **64**, 1139-1142 (2001)



Japanese Sea mussel *Mytilus coruscus*



Takashi Maoka and Takao Matsuno, Isolation and structural elucidation of three new acetylenic carotenoids from the Japanese sea mussel *Mytilus coruscus*, *Nippon Suisan Gakkaishi*, **54**, 1443-1447 (1988).



Takashi Maoka, A New Apocarotenoid From Marine Shellfish, *J. Nat. Prod.*, **60**, 616-617 (1997).

Scallop *Patinopecten yessoensis*



S. A. Campbell, A.K. Malla,us, E.S. Weight, B. C. L. Weedon, Pectenoxanthin, cynthiaxanthin and a new acetylenic carotenoid pectenolone, Chem. Commun. 941 –942 (1967).

Ark Shell *Scapharca broughtonii*



Takao Matsuno and Takashi Maoka, Isolation of diatoxanthin, pectenoxanthin, pectenolone, and a new acetylenic carotenoid 3,4,3'-trihydroxy-7',8'-didehydro- -carotene from the arkshell and related three species of bivalves, Bull. Jap. Soc. Sci. Fish., 47, 495-499 (1981).